

# PATENT COOPERATION TREATY

EO/US  
PCT/AU93/00230

**PCT**

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark  
Office  
Washington, D.C.

in its capacity as elected Office

Date of mailing:

25 November 1993 (25.11.93)

International application No.:

PCT/AU93/00230

Applicant's or agent's file reference:

469942

International filing date:

19 May 1993 (19.05.93)

Priority date:

19 May 1992 (19.05.92)

Applicant:

KELLY, Graham, Edmund

1. The designated Office is hereby notified of its election made:



in the demand filed with the International preliminary Examining Authority on:

11 October 1993 (11.10.93)



in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was



was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO  
34, chemin des Colombettes  
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer:

J. Zahra

Telephone No.: (41-22) 730.91.11

# PATENT COOPERATION TREATY

## PCT


### INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rules 70)

Applicant's or agents file reference 469942	<b>FOR FURTHER ACTION</b>	See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416).
International application No.  <b>PCT/AU 93/00230</b>	International filing date  19 May 1993 (19.05.93)	Priority Date  19 May 1992 (19.05.92)
International Patent Classification (IPC) or national classification and IPC <b>Int. Cl. <sup>5</sup> A61K 37/24; 35/78</b>		
Applicant <b>KELLY, Graham Edmund</b>		

REC'D 28 JUN 1994  
 WIPO PCT

1.	This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2.	This REPORT consists of a total of 4 sheets, including this cover sheet.  <input type="checkbox"/> This report is also accompanied by ANNEXES, i.e., sheet of description, claim and/or drawings amended during international preliminary examination and/or containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).  These annexes consist of a total of    sheets.
3.	This report contains indications relating to the following items:  <div style="margin-left: 20px;">           I    <input checked="" type="checkbox"/> Basis of the report             II   <input type="checkbox"/> Priority             III <input type="checkbox"/> Non-establishment of opinion with regard to novelty, inventive step and industrial applicability             IV   <input type="checkbox"/> Lack of unity of invention             V    <input checked="" type="checkbox"/> Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement             VI   <input type="checkbox"/> Certain documents cited             VII <input type="checkbox"/> Certain defects in the international application             VIII <input checked="" type="checkbox"/> Certain observations on the international application         </div>

Date of submission of the demand 11 October 1993 (11.10.93)	Date of completion of this report
Name and mailing address of the IPEA/AU AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION PO BOX 200 WODEN ACT 2606 AUSTRALIA  Facsimile No. (06) 2853929	Authorized officer  <div style="text-align: center;">   <b>J.P. PULVIRENTI</b> </div> Telephone No. (06) 2832261

**I. Basis of the report**

1. This report has been drawn on the basis of *(Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.)*:

- ☐ the international application as originally filed.
- ☒ the description,                      pages 1-20, as originally filed,  
   pages ..., as filed with the demand,  
   pages ..., filed with the letter of ...,  
   pages ..., filed with the letter of ...,
- ☒ the claims,                              Nos. 11-20, as originally filed,  
   Nos. 1-10, as amended under Article 19,  
   Nos. ..., filed with the demand,  
   Nos. ..., filed with the letter of ...,  
   Nos. ..., filed with the letter of ... ,
- ☐ the drawings,                           sheets/fig ..., as originally filed,  
   sheets/fig ..., filed with the demand,  
   sheets/fig ..., filed with the letter of ...,  
   sheets/fig ..., filed with the letter of ...,

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages ...
- ☐ the claims, Nos. ...
- ☐ the drawings, sheets/fig ...

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).

4. Additional observations, if necessary:

**V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

**1. STATEMENT**

Novelty (N)	Claims 1-20	YES
	Claims	NO
Inventive Step (IS)	Claims	YES
	Claims 1-20	NO
Industrial Applicability (IA)	Claims 1-20	YES
	Claims	NO

**2. Citations and explanations**

**INVENTIVE STEP (IS) Claims 1-20**

- (a) EP,A,0135172 (TAKEDA CHEMICAL INDUSTRIES LTD) 27 March 1985 (27.03.85)
- (b) Patent Abstracts of Japan C-452, page 118, JP,A,62-106016 (YAMANOUCHI PHARMACEUT CO LTD) 16 May 1987 (16.05.87)
- (c) Patent Abstracts of Japan C-722, page 98, JP,A,2-67218 (NAGAKURA SEIYAKU K.K.) 7 March 1990 (07.03.90)
- (d) Chemical Abstracts, volume 115, No. 8, issued 26 August 1991 (26.08.91) (Columbus Ohio) USA. LIU, Y. et al. "Effects of solid dispersion of daidzein on the blood pressure of spontaneously hypertensive rats", page 406, abstract no. 78763p, Shenyang Yaoxueyuan Xuebao, 1991, 8(2) 105-8 (Ch.)

These citations all disclose the use of the claimed phyto-oestrogens in treatments to improve human health. It would be obvious to a person skilled in the art to combine the claimed compounds to improve human health. The claimed phyto-oestrogens are all individually known as agents which improve human health. There has been no surprising or unexpected result demonstrated in using these compounds together as a health supplement which is beyond the aggregate of the known health improving properties of these phyto-oestrogens.

**VIII. Certain observations on the international application**


The following observations on the clarity of the claims, descriptions, and drawings or on the question whether the claims are fully supported by the description, are made:

- (a) Claim 1 lacks clarity as the ambit of "health supplement" and "health supplementary amount" is unclear. Similarly in claim 10, the ambit of "method of improving the health" and "health supplementing amount" is unclear. The known uses of the claimed compounds (e.g. osteoporosis treatment) are treatments to improve a person's health, and it is not clear how these terms serve to distinguish the invention from the prior art.
- (b) Claims 1 and 10 lack clarity as the meaning of naturally occurring phyto-oestrogen is unclear. The phyto-oestrogens defined in these claims inherently occur in nature, and it is not clear whether this term includes all these phyto-oestrogens, or only those extracted from natural sources (e.g. red clover), as exemplified in the specification.

If all of these natural products are included then these claims lack novelty. If they are not included then the distinction between these natural sources of the extracts, and the final phyto-oestrogen composition is not clearly defined.

- (c) Note: Whilst the ISR and this report covered the methods of claims 10-20, other member countries of the PCT may regard these claims as being excluded subject matter under Art. 17(2)(a)(i) [Rule 39.1].

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate of the relevant passages	Relevant to Claim No.
X	Patent Abstracts of Japan C-452, page 118, JP,A,62-106016 (YAMANOUCHI PHARMACEUT CO. LTD) 16 May 1987 (16.05.87) see entire abstract	1-20
X	Patent Abstracts of Japan C-722, page 98, JP,A,2-67218, (NAGAKURA SEIYAKU K.K.) 7 March 1990 (07.03.90) see entire abstract particularly formula I	1-20
X	Chemical Abstracts, volume 115, No. 8, issued 26 August 1991 (26.08.91) (Columbus, Ohio) USA. LIU, Y. et. al. "Effects of solid dispersion of daidzein on the blood pressure of spontaneously hypertensive rats", page 406, abstract no. 78763p, Shenyang Yaoxueyuan Xuebao 1991, 8(2) 105-8 (Ch.) see entire abstract	1-20
A	EP,A,0136569 (TAKEDA CHEMICAL INDUSTRIES LTD) 10 April 1985 (10.04.85) see pages 1, 6-7	1-20
A	GB,A,1482238 (CHINOIN GYOGYSZER ES VEGYESZETI TERMEKEK GYARA RT), 10 August 1977 (10.08.77) see pages 1 and 4	1-20

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int. Cl. <sup>5</sup> A61K 37/24, 35/78  According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b>  Minimum documentation searched (classification system followed by classification symbols) IPC A61K 37/24, 35/78  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched AU: IPC as above				
Electronic data base consulted during the international search (name of data base, and where practicable, search terms used) DERWENT: GENISTEIN: OR DAIDZEIN: OR BIOCHANIN: OR FORMONONETIN: JAPIO: GENISTEIN: OR DAIDZEIN: OR BIOCHANIN: OR FORMONONETIN:				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.		
X	EP,A,0135172 (TAKEDA CHEMICAL INDUSTRIES LTD) 27 March 1985 (27.03.85) see pages 1, 8 and 9	1-20		
X	EP,A,0129667 (TAKEDA CHEMICAL INDUSTRIES LTD) 2 January 1985 (02.01.85) see page 5, page 7 lines 17-35	1-20		
X	Patent Abstract of Japan C-757, page 17, JP,A,2-160722 (NIPPON OIL AND FATS CO LTD) 20 June 1990 (20.06.90) see entire abstract	1-20		
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.</span> <span><input checked="" type="checkbox"/> See patent family annex.</span> </div>				
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top;"> <p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </td> <td style="width: 50%; vertical-align: top;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p> </td> </tr> </table>			<p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p>
<p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p>			
Date of the actual completion of the international search 3 August 1993 (03.08.93)		Date of mailing of the international search report 19 AUG 1993 (19.08.93)		
Name and mailing address of the ISA/AU  AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION PO BOX 200 WODEN ACT 2606 AUSTRALIA  Facsimile No. 06 2853929		Authorized officer    <b>J. PULVIRENTI</b>  Telephone No. (06) 2832253		

**A. CLASSIFICATION OF SUBJECT MATTER**Int. Cl.<sup>5</sup> A61K 37/24, 35/78

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC A61K 37/24, 35/78

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

AU: IPC as above

Electronic data base consulted during the international search (name of data base, and where practicable, search terms used)

DERWENT: GENISTEIN: OR DAIDZEIN: OR BIOCHANIN: OR FORMONONETIN:

JAPIO: GENISTEIN: OR DAIDZEIN: OR BIOCHANIN: OR FORMONONETIN:

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	EP,A,0135172 (TAKEDA CHEMICAL INDUSTRIES LTD) 27 March 1985 (27.03.85) see pages 1, 8 and 9	1-20
X	EP,A,0129667 (TAKEDA CHEMICAL INDUSTRIES LTD) 2 January 1985 (02.01.85) see page 5, page 7 lines 17-35	1-20
X	Patent Abstract of Japan C-757, page 17, JP,A,2-160722 (NIPPON OIL AND FATS CO LTD) 20 June 1990 (20.06.90) see entire abstract	1-20

☒ Further documents are listed  
in the continuation of Box C.☒ See patent family annex.**\* Special categories of cited documents :**

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search  
3 August 1993 (03.08.93)

Date of mailing of the international search report  
19 AUG 1993 (19.08.93)

Name and mailing address of the ISA/AU

AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION  
PO BOX 200  
WODEN ACT 2606  
AUSTRALIA

Facsimile No. 06 2853929

Authorized officer

J. PULVIRENTI

Telephone No. (06) 2832253



C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate of the relevant passages	Relevant to Claim No.
X	Patent Abstracts of Japan C-452, page 118, JP,A,62-106016 (YAMANOUCHI PHARMACEUT. CO. LTD) 16 May 1987 (16.05.87) see entire abstract	1-20
X	Patent Abstracts of Japan C-722, page 98, JP,A,2-67218, (NAGAKURA SEIYAKU K.K.) 7 March 1990 (07.03.90) see entire abstract particularly formula I	1-20
X	Chemical Abstracts, volume 115, No. 8, issued 26 August 1991 (26.08.91) (Columbus, Ohio) USA. LIU, Y. et. al. "Effects of solid dispersion of daidzein on the blood pressure of spontaneously hypertensive rats", page 406, abstract no. 78763p, Shenyang Yaoxueyuan Xuebao 1991, 8(2) 105-8 (Ch.) see entire abstract	1-20
A	EP,A,0136569 (TAKEDA CHEMICAL INDUSTRIES LTD) 10 April 1985 (10.04.85) see pages 1, 6-7	1-20
A	GB,A,1482238 (CHINOIN GYOGYSZER ES VEGYESZETI TERMEKEK GYARA RT), 10 August 1977 (10.08.77) see pages 1 and 4	1-20

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member			
EP	135172	DE A1 3430799 JP A2 60048924	IT A 8467840	IT	A 1179067
EP	129667	DE A 3415394 JP A 59199630	IT A 8467428	IT	A 1208776
EP	136569	HU A 36111	JP A 60054379		
GB	1482238	AR A 205818 AT A 6908/77 CH A 611289 DE A 2518986 FI B 63568 FR B 2269340	AT A 3320/75 AT B 351919 CS P 183794 DK A 1913/75 FI C 63568 HU P 168753	AT B 344167 AU A 80723/75 DD Z 119587 FI A 751272 FR A 2269340 IL A 47196	
END OF ANNEX					

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member			
EP	135172	DE A1 3430799 JP A2 60048924	IT A 8467840	IT	A 1179067
EP	129667	DE A 3415394 JP A 59199630	IT A 8467428	IT	A 1208776
EP	136569	HU A 36111	JP A 60054379		
GB	1482238	AR A 205818 AT A 6908/77 CH A 611289 DE A 2518986 FI B 63568 FR B 2269340	AT A 3320/75 AT B 351919 CS P 183794 DK A 1913/75 FI C 63568 HU P 168753	AT B 344167 AU A 80723/75 DD Z 119587 FI A 751272 FR A 2269340 IL A 47196	
END OF ANNEX					

## PATENT COOPERATION TREATY

PCT

NOTIFICATION CONCERNING  
DOCUMENT TRANSMITTED

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark  
Office  
Washington, D.C.

in its capacity as elected Office

Date of mailing:

30 June 1994 (30.06.94)

International application No.:

PCT/AU93/00230

International filing date:

19 May 1993 (19.05.93)

Applicant:

KELLY, Graham, Edmund

The International Bureau transmits herewith the following documents and number thereof:

\_\_\_\_\_ copy of the international preliminary examination report and annexes (Article 36(3)(a))

The International Bureau of WIPO  
34, chemin des Colombettes  
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorised officer:

M. Abidine

Telephone No.: (41-22) 730.91.11

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

/338567



55 Rec'd PCT/PTC

21 NOV 1994

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 5 :</b> <b>A61K 37/24, 35/78</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/23069</b> <b>(43) International Publication Date:</b> 25 November 1993 (25.11.93)
<b>(21) International Application Number:</b> PCT/AU93/00230 <b>(22) International Filing Date:</b> 19 May 1993 (19.05.93)  <b>(30) Priority data:</b> PL 2511 19 May 1992 (19.05.92) AU  <b>(71)(72) Applicant and Inventor:</b> KELLY, Graham, Edmund [AU/AU]; 112 Kameruka Road, Northbridge, NSW 2063 (AU).  <b>(74) Agent:</b> GORDON, Glen, Howard; Davies Collison Cave, Level 10, 10 Barrack Street, Sydney, NSW 2000 (AU).  <b>(81) Designated States:</b> AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).		<b>Published</b> <i>With international search report.</i> <i>With amended claims and statement.</i>
<b>(54) Title:</b> HEALTH SUPPLEMENTS CONTAINING PHYTO-OESTROGENS, ANALOGUES OR METABOLITES THEREOF  <b>(57) Abstract</b>  Compositions enriched with natural phyto-oestrogens or analogues thereof selected from Genistein, Daidzein, Formononetin and Biochanin A. These may be used as food additives, tablets or capsules for promoting health in cases of cancer, pre-menstrual syndrome, menopause or hypercholesterolaemia.		

HEALTH SUPPLEMENTS CONTAINING PHYTO-OESTROGENS, ANALOGUES OR METABOLITES THEREOF

TECHNICAL FIELD

This invention relates to natural products containing phyto-oestrogens, or phyto-oestrogen metabolites, which have various beneficial physiological effects in man, and which have a variety of uses, such as to promote good health and as a dietary additive, for example.

BACKGROUND ART

The particular product in accordance with the invention is an extract of certain plants with the particular purpose of enrichment for phyto-oestrogens, both in their natural state and their closely related derivatives and metabolites.

Plants which are used as foodstuffs or medicinal herbs contain a wide variety of chemicals which are assimilated into the body following ingestion. Some of these chemicals are important nutrients for man and animals (e.g. fats, carbohydrates, proteins, vitamins, minerals) while others have none, or little or no known nutritional value. The phyto-oestrogens hitherto have fallen into this latter category of no known nutritional value.

There are 3 principal classes of phyto-oestrogens, viz. isoflavones, lignans, and coumestans. The isoflavones are thought to have a broad range of biological functions in plants, although these are poorly understood. However, two particular functions are recognised - (a) as phytoalexin or stressor chemicals which are secreted by the plant in response to attack by parasites such as insects, fungi, viruses, etc and which display activity against these parasites, and (b) chemicals which encourage colonisation of nitrogen-fixing bacteria on the roots of legumes. The biological functions in plants of the lignans and coumestans is not generally understood.

The different types of phyto-oestrogens are as follows.

Type 1 phyto-oestrogens - (isoflavones)

Isoflavones appear to be widely distributed in the plant kingdom and over 700 different isoflavones are described. However, the isoflavones which display oestrogenic activity belong to a small sub-group and are restricted almost exclusively to the *Leguminosae* family. The known oestrogenic isoflavones are daidzein, formononetin, genistein and biochanin A. In

common human foodstuffs such as soya, chickpeas, lentils and beans, the total levels of the oestrogenic isoflavones range between about 40 and 300 mg per 100 g dry weight.

In the raw plant material, isoflavones occur principally as glycosides. Following ingestion by man and animals, the glycoside moiety is hydrolysed free by a combination of gastric acid hydrolysis and fermentation by intestinal bacteria. Some of the isoflavones in the aglucone form are absorbed directly and circulate in the blood, while the remainder are metabolised by intestinal fermentation to a variety of compounds which are also absorbed. The absorbed isoflavones and their metabolites appear to undergo little or no further metabolism in the body, being readily transported in the bloodstream, and ultimately being excreted in the urine.

#### Type 2 phyto-oestrogens (lignans).

Lignans are widely distributed in the plant kingdom. Over one hundred lignans are described and they are reported in common human foodstuffs such as cereals, fruits and vegetables. Oilseeds such as flax (linseed) have the highest known levels at 20-60 mg/100 g dry weight, while cereals and legumes have much lower levels at 0.3-0.6 mg/100 g, and vegetables even lower levels at 0.1-0.2 mg/100 g. The most common lignan described is metairesinol. Dietary lignans also appear to be metabolised fairly efficiently within the gut by bacterial fermentation, yielding metabolites such as enterodiol and enterolactone which are absorbed into the bloodstream and excreted in the urine.

#### Type 3 phyto-oestrogens (coumestans).

Compared to isoflavones and lignans, oestrogenic coumestans appear to have a relatively restricted distribution in plants and generally occur at much lower levels. Alfalfa, ladino clover and some other fodder crops such as barrel medic may have significant levels and have been reported to cause reproductive dysfunction in grazing animals. In the human diet, the important sources of coumestans are sprouts of soya and alfalfa where levels up to 7 mg/100g dry weight are reported. Whole soyabeans and other common foodstuff legumes contain levels of approx. 0.12 mg/100 g dry weight and most of that is concentrated in the seed hull which commonly is removed in the preparation of human foodstuffs.

#### Type 4 phyto-oestrogens (oestrogens).

These are compounds closely related to animal oestrogens such as oestrone, oestradiol and

3

oestriol. These have been described in plants such as liquorice, apple, French bean, pomegranate and date palm. Little is known of the metabolism and biological significance of these chemicals in humans and animals.

The full range of biological effects in animals of these dietary phyto-oestrogens has received only recent study. A primary effect appears to be associated with their close structural relationship to naturally-occurring oestrogens which allows the phyto-oestrogens to mimic the effects of the endogenous oestrogens. The known biological effects of phyto-oestrogens can be summarised thus:

- In vitro*
- (a) bind to both cytoplasmic and nuclear membrane (Type II) oestrogen receptors on human tissues;
  - (b) strongly compete with oestrogens for oestrogen receptors, but only weakly stimulate those receptors;
  - (c) strongly stimulate the production of sex hormone-binding globulin (SHBG) from human cells;
- In vivo*
- (d) weakly oestrogenic in animals;
  - (e) competitively-inhibit the response of tissue to oestrogens.

The three major types of phyto-oestrogens appear to act at the cellular level in a similar manner, that is through interaction with cell surface oestrogen receptors. In the body, naturally-occurring oestrogens circulating in the blood largely exert their activity by interaction with oestrogen receptors on cell surfaces; such interactions then triggering a particular biological function of that particular cell. Phyto-oestrogens are able to bind to those oestrogen receptors because the structure of these compounds so closely resembles the endogenous oestrogens, but unlike the animal oestrogens, phyto-oestrogens only weakly activate the oestrogen receptor.

As a result of phyto-oestrogens and endogenous oestrogens competing for the oestrogen-binding sites on cells, the more weakly oestrogenic phyto-oestrogens can be considered to have an anti-oestrogenic effect. This phenomenon is known as competitive-inhibition, by which is meant that the biological effect of an active substance is impaired by the competitive

4



binding to a target receptor of a similar but less active compound.

Thus a primary biological effect of phyto-oestrogens is held to be competitive inhibition of endogenous oestrogens. However, another more direct effect is the stimulation of synthesis of SHBG in the liver, as occurs with orally administered synthetic steroidal oestrogens. High levels of dietary phyto-oestrogens are thought to be responsible for the higher SHBG levels seen in vegetarians and in cultures maintaining traditional (high legume-containing) diets.


At high levels, dietary phyto-oestrogens can have profound physiological effects. An example of this is sheep and cattle grazing pastures containing a high proportion of subterranean clover or red clover which can contain levels of phyto-oestrogens as high as 5% of the dry weight of the plant. As a result of the competitively-inhibitory effect of the dietary phyto-oestrogens on endogenous oestrogen function in the hypothalamus, male and female sheep and cows can develop androgenic symptoms.

Such high dietary levels of phyto-oestrogens, however, are rare. It is far more common that most animal and human diets contain low to moderate levels of phyto-oestrogens, and there is growing epidemiological evidence that such levels have a beneficial effect on human health.

In most traditional human diets in developing countries, the principal phyto-oestrogens consumed are isoflavones because of the generally high reliance on legumes (also known as pulses) as a source of protein. The general consumption rates (g/day/person) for legumes for different regions currently are approximately: Japan (50-90), India (40-80), South America (30-70), North Africa (40-50), Central/Southern Africa (20-50) and Southern Mediterranean (30-60). Legumes also are a source of lignans and, to a much lesser extent, coumestans, and the additional cereal and vegetables in the diet would also boost the lignan intake. However, the isoflavone intake in these traditional cultures with high legume consumption would typically be much in excess of either lignan or coumestan intake.

The major types of legumes used in traditional diets include soya, chickpeas, lentils, ground nuts, beans (e.g. broad, haricot, kidney, lima, navy), and grams (bengal, horse and green).

In Western, developed countries, the daily intake of dietary phyto-oestrogens generally is



negligible to low. In Western Europe, North America and Australasia, legumes were a major source of protein for the majority of the populations up to the end of the 19th century. From that time, legume consumption has declined significantly, being replaced in the diet with protein of animal origin. Average legume consumption in these regions currently is between 5-15 g/day/person with a significant proportion of the population ingesting little to no legumes or other phyto-oestrogen containing foods on a regular basis. Moreover, the types of legumes consumed in these regions (e.g. garden peas, French beans) have a typically lower isoflavone content than legumes such as soya and chick peas.

Based on typical consumption rates and types of foodstuffs consumed, the typical phyto-oestrogen intake (mg/day) for different regions can be calculated approximately as

TCox

	<u>Isoflavones</u>	<u>Lignans</u>	<u>Coumestans</u>
Japan	50-300	2-5	0.5
Australia	2-25	1-5	0.2

Thus it can be seen that regions which have maintained traditional diets have a higher average daily intake of phyto-oestrogens, particularly isoflavones, compared to western countries. People in communities such as Japan or developing countries with high legume intake excrete substantially higher phyto-oestrogen metabolites in their urine compared to people in Western countries. Within the latter, vegetarians also excrete higher phyto-oestrogen metabolite levels than do those consuming a more typical, omnivorous Western diet.

The presence of relatively large amounts of phyto-oestrogen metabolites in urine serves to highlight their potential biological significance. It has been shown that total urinary excretion of isoflavones and their active metabolites in people consuming moderate amounts of legumes is greatly in excess (up to 10,000 x) of steroidal oestrogen levels. So that while the oestrogenicity of isoflavones to oestrogen receptors is only about 1% that of endogenous oestrogens, this weaker effect is off-set by the much higher blood levels of the isoflavones.

It is known that legumes have formed an important part of the human diet over the past 20,000-30,000 years. It therefore follows that human metabolism has evolved over at least this period in the presence of relatively large levels of dietary phyto-oestrogens, particularly

isoflavones. Given the known biological effects of phyto-oestrogens, it also follows that endogenous oestrogen metabolism and function has evolved in the face of significant competitive inhibiting effects of phyto-oestrogens. It has been speculated that the presence of significant dietary levels of phyto-oestrogens in recent human evolution has led to a degree of adaption by tissues responsive to reproductive hormones to these dietary components. That is, both the rate of production and/or the function of endogenous oestrogens may be either dependent upon or influenced by the presence of phyto-oestrogens in the body. It follows therefore that a relative deficiency of dietary phyto-oestrogens could be expected to lead to an imbalance of endogenous oestrogen metabolism.

There is increasing interest in the likely contribution of a relative deficiency of dietary phyto-oestrogens to the development of the so-called "Western diseases", especially cancer of the breast, benign (cystic) breast disease, cancer of the uterus, cancer of the prostate, cancer of the bowel, pre-menstrual syndrome, menopausal syndrome, and atherosclerosis. All of these diseases are associated to a greater or lesser extent to oestrogen metabolism, and oestrogen function is either known or is suspected to play a role in their aetiology and/or pathogenesis.

Each of these diseases occurs at much higher incidence in Western, developed countries than it does in developing communities. Moreover, it is thought that in Western communities, the incidences of each have risen over the past century. It is also generally held, that of all the environmental factors likely to be contributing to this phenomenon, diet is the principal factor. Of those dietary components with the potential to influence the aetiology of oestrogen-related disease, there is a growing awareness that phyto-oestrogens may have important potential.

The beneficial effects of phyto-oestrogens on human health are thought to derive from at least two principal function, those being (i) competitive-inhibition of the function of endogenous oestrogens, and (ii) the stimulation of production of SHBG. SHBG plays an important role in primates in binding and transporting the reproductive hormones (oestrogens, androgens) in blood so that the availability of reproductive hormones is regulated to a large degree by SHBG levels. Higher SHBG levels are considered beneficial in leading to a reduction in both blood levels of unbound (and unregulated) reproductive hormones and metabolic clearance rates of the hormones. Although isoflavones are potent stimulators of SHBG synthesis, they only weakly bind to SHBG, so that the increased SHBG levels resulting from the dietary

7

isoflavones are largely available for binding to endogenous oestrogens.

In terms of directly identifying the beneficial effects of phyto-oestrogens in amelioration of any or all of the "Western diseases", there are only two examples. In one example, the diets of women, with menopausal syndrome were supplemented with foodstuffs (soya, linseed, red clover) high in phyto-oestrogens, and an alleviation of menopausal symptoms to an extent similar to that obtained with replacement therapy with synthetic oestrogens was achieved; that effect was ascribed to the phyto-oestrogen content of the supplement. In the other example, legumes such as soya and various pulses have been shown to have a hypocholesterolaemic effect in humans; this effect has not been ascribed to phyto-oestrogens, although purified isoflavones do have a hypocholesterolaemic effect in animals with artificially-induced hypercholesterolaemia.

In summary, it could reasonably be deduced that the inclusion of greater levels of foodstuffs high in phyto-oestrogens in the standard diets of men and women in developed countries could be expected to redress a general imbalance of endogenous reproductive hormone metabolism, thereby reducing the predisposition of those communities to the above diseases. While there are various types of phyto-oestrogens which may be suitable to this end, the large discrepancy in isoflavone consumption between communities with Western and traditional diets suggest that foodstuffs with high isoflavone content are of prime interest.

However it is unrealistic to expect that public education programmes would readily convert communities in developed countries from a diet where the protein content is predominantly animal-derived, to one where the protein is predominantly legume-derived. Moreover, the legumes which are commonly consumed in developed countries are relatively poor sources of phyto-oestrogens and the general acceptance in the community of less well-known legumes with higher phyto-oestrogen content would be necessarily a slow process. Also, the highly variable levels of phyto-oestrogens in foodstuffs relating to plant strain type, degree of plant maturity, and climatic and other environmental conditions suggests that the supply of an assured amount of phyto-oestrogens through the use of whole foodstuffs may be difficult.

An alternative strategy is to make available either (i) phyto-oestrogens in a purified form, or (ii) foodstuffs which are enriched for phyto-oestrogens. In this way, the phyto-oestrogen could

be added to the diet in a convenient form as a supplement without requiring any substantive change to the diet.


#### DISCLOSURE OF INVENTION

The present invention concerns a health supplement specifically enriched for isoflavones selected from genistein, daidzein, formononetin and biochanin A, or their natural glycoside form, or their analogues, in sufficient amounts to improve the health of a human.

Preferably the supplement contains an excipient, a diluent, a carrier or the like, or else the supplement is mixed with food or can be consumed directly. It is also preferred that foodstuffs, are readily available, have no known toxic components, and are rich sources of isoflavones; such foodstuffs preferably being red clover or soya. It is also preferred that the ratio of genistein and/or its methylated derivative biochanin A to daidzein and/or its methylated derivative formononetin is between 1:2 to 2:1. Other plant components with oestrogenic activity including lignans, coumestans and flavones may also be present in the extract, but it is held that these are of secondary importance to the predominant isoflavones. The term phyto-oestrogens is used hereafter to indicate a predominance of isoflavones with lesser amounts of lignans, coumestans and flavones.


The invention also concerns a method of improving the health of a human by administering to the human a sufficient amount of phyto-oestrogen. Ideally, the phyto-oestrogen is administered regularly on a daily basis over a sufficient period such as at least a month. The health conditions which may be prevented or ameliorated include cancer of the breast, cancer of the prostate, cancer of the uterus, cancer of the bowel, benign (or cystic) breast disease, pre-menstrual syndrome (also known as pre-menstrual tension), or adverse symptoms associated with menopause in women. The method and supplement in accordance with the invention also improves the health of a human having elevated levels of blood cholesterol. The product also is useful in avoiding or ameliorating cancer in persons. The symptoms produced by these conditions and the general well-being is also improved by the use of these supplements.

The phyto-oestrogen in accordance with the invention may be obtained from a number of different sources. Preferably the phyto-oestrogens are extracted from a clover such as red clover or subterranean clover or from soya which contain high levels of phyto-oestrogens.



However, any source rich in phyto-oestrogens may be used instead, if desired.

Various different isoflavones have been identified from these sources - they are principally genistein, biochanin A, daidzein, formononetin and glycitein. In plants these compounds occur principally in a glycoside form bound to sugars such as glucose, with smaller amounts present as the aglucone forms. The formulae of the isoflavones are:



The structure of biochanin A is the same as for genistein but with a 4'-methoxy group, and similarly formononetin has the same structure as daidzein, but with a 4'-methoxy group.

Following ingestion by humans, the glycosidic isoflavones are hydrolysed to the aglucone form and biochanin A and formononetin are demethylated by bacterial fermentation to genistein and daidzein respectively. A small proportion of these free isoflavones are absorbed directly from the bowel and circulate in the blood. The bulk of the isoflavones, however, remain in the bowel and undergo fermentation to form various metabolites which also are absorbed into the bloodstream. The principal metabolites which have been identified are equol and O-desmethylangolensin.

*In vitro* and *in vivo* studies have indicated that genistein, biochanin A, equol, daidzein, formononetin all have oestrogenic activity in descending order. O-desmethylangolensin is only very weakly oestrogenic and glycitein is non-oestrogenic.

In animal and *in vitro* studies, genistein has been shown to have greater oestrogenic/anti-oestrogenic activity and SHBG-stimulating capacity than the other isoflavones or their metabolites (approximately 10 times that of daidzein and formononetin). However, the full range of biological effects of the different isoflavones have yet to be fully determined, and in particular their relative efficacies in the different biological effects such as oestrogenicity,

10

hypcholesterolaemia, anti-angiogenesis, anti-oxidation, anti-carcinogenesis for example are not yet fully known.

It is thought that because the methyl forms (biochanin A and formononetin) ultimately are largely demethylated to their principals, genistein and daidzein, with improved biological efficacy, then it is unimportant whether the isoflavones are present in the claimed product in the methylated or demethylated forms.

Given that the relative biological importance of the two isoflavone groups (being genistein and daidzein) to human health remains unclear, and that each might indeed have different importance, plus the fact that both isoflavones are present in the diet in approximately equal proportions, then it is prudent that both isoflavones be present in the claimed product in approximately equal proportions.

Any leguminous plants such as detailed here could be used as sources of phyto-oestrogens (principally isoflavones with lesser amounts of lignans and coumestans): Indian liquorice (*Abrus precatorius*); various species of *Acacia* spp. including, *A. aneura*, *A. cibaria*, *A. longifolia*, and *A. oswaldii*; ground nut (*Apios tuberosa*); ground pea (*Arachis hypogaea*); milk vetch (*Astragalus edulis*); marama bean (*Bauhinia esculenta*); sword bean (*Cajanus cajan indicus*); jack bean (*Canavalia ensiformis*); sword bean (*Canavalia gladiata*); seaside sword bean (*Canavalia rosea*); various *Cassia* spp. including *C. floribunda*, *C. laevigata*, and *C. occidentalis*; carobbean (*Ceratonia siliqua*); chick pea (*Cicer arietinum*); yebnut (*Cordeauxia edulis*); various *Crotalaria* spp. including *C. laburnifolia*, and *C. pallida*; cluster bean (*Cyamopsis psoralioides*); tallow tree (*Detarium senegalense*); sword bean (*Entada scandens*); balu (*Erythrina edulis*); soyabean (*Glycine max*); inga (*Inga edulis*); Polynesian chestnut (*Inocarpus fagifer*); hyacinth bean (*Lablab purpureus*); grass pea or Indian vetch (*Lathyrus sativus*); cyprus vetch (*Lathyrus ochrus*); lentil (*Lens culinaris*); jumping bean (*Leucaena leucocephala*); various *Lupinus* spp. including *L. albus*, *L. luteus*, *L. angustifolium*, *L. mutabilis*, and *L. cosentinii*; ground bean (*Macrotyloma geocarpa*); horse gram (*Macrotyloma uniflorum*); alfalfa (*Medicago sativa*); velvet bean (*Mucuna pruriens*); yam beans (*Pachyrhizus erosus*, *P. tuberosus*); African locust bean (*Parkia clappertoniana*); *Parkia speciosa*; oil bean tree (*Pentaclethra macrophylla*); various *Phaseolus* spp. including *P. acutifolius*, *P. vulgaris*, *P. luntus*, *P. coccineus*, *P. adenathus*, *P. angulris*, *P. aureus*, *P. calcaratus*, *P. mungo*, and *P.*

*polystachyus*; garden pea (*Pisum sativum*); djenko bean (*Pithecolobium lobatum*); mesquite (various *Prosopis* spp.); goa bean (*Psophocarpus scandens*, *P. tetragonolobus*); various *Psoralea* spp.; *Sesbania bispinosa*; yam bean (*Sphenostylis stenocarpa*); tamarind (*Tamarindus indica*); fenugreek (*Trigonella foenum-graecum*); vetches (various *Vicia* spp. including *V. sativa*, *V. atropurpurea*, *V. ervilia*, and *V. monantha*); broad bean (*Vicia faba*); black gram (*Vigna mungo*); various *Vigna* spp. including *V. radiata*, *V. aconitifolia*, *V. adanatha*, *V. angularis*, *V. tribolata*, *V. umbelata*, and *V. unguiculata*; and, earth pea (*Voandzeia subterranea*).

The ideal sources of phyto-oestrogens for preparation of a supplement in accordance with the invention are preferably those which (i) are readily available, (ii) are relatively inexpensive, (iii) are readily and economically processed so as to yield the extract, (iv) have a high isoflavone content so as to provide high yields, and (v) have no known toxic components requiring selective removal or inactivation.

Certain clovers, such as red clover (*T. pratense*) and subterranean clover (*T. subterranean*) are the preferred sources. On a dry weight basis, these clovers contain the highest amounts of oestrogenic isoflavones of all legumes tested to date with levels of 3-5 g% (*T. subterranean*) and 1-3 g% (*T. pratense*). In comparison, soya flour has a level of 0.15-0.30 g%, lentils (0.08-0.12 g%), chick peas (0.07-0.13 g%), and garden peas (0.02-0.03 g%). Thus it can be seen that clovers contain approximately at least 10-30 times by weight the isoflavone content of other commonly available, human leguminous foodstuffs meaning that for manufacturing purposes, the yield of isoflavones per unit weight of plant material is many times greater from clover than from other legumes.

Red clover and subterranean clover also are common fodder crops and are readily grown and are widely available. Clovers also are comparatively cheaper (\$200/tonne) than crops such as soya and lentils (\$500/tonne).

With clovers, the isoflavones are recovered from the leaf rather than from the seed in the case of soya, beans, nuts and grams. This provides a substantially higher yield of isoflavones per unit area of pasture for clovers compared to other legumes because of the greater leaf matter compared to seed matter recovered per plant.



Clovers also have an extended growing season, and faster growth rates compared to those legumes such as soya, lentils or chick peas where the seed is the end-product. Clover can be cropped for its leaf content repeatedly over a single growing season. An additional benefit of this is that as phyto-alexins, the isoflavone content increases in response to the stress of cropping.

Thus it can be seen that in clovers versus other legumes provide a combination of (a) higher isoflavone content per dry weight of plant, (b) a higher yield of dry matter containing isoflavones per plant, and (c) a higher yield of dry matter per hectare.

An additional feature of clovers is that there are wide varieties of cultivars with widely differing isoflavone levels and types. This allows blending of different cultivars to achieve the desired ratio of the different isoflavones, although it is equally possible to use a single cultivar which provides the desired ratio.

Other legumes such as soyabean flour may be used for enrichment of phyto-oestrogens but the substantially poorer (approx. 10%) yield of isoflavones compared to clovers means that the manufacturing costs are substantially greater and there is substantially greater amounts of waste products which requires disposal or further treatment for re-use as a foodstuff. An alternative, however, to the use of whole soya for this purpose, is to use the hull and hypocotyl (or germ) of the whole soyabean. The hull and hypocotyl represent only a small proportion by weight (8% and 2% respectively) of the intact bean. However, the coumestrol content of soya is concentrated in the hull, and the daidzein content of soya is concentrated in the hypocotyl. The two cotyledons which comprise the bulk of the soyabean (90% by weight) contain the bulk of the genistein content of soya. During standard processing of soyabeans, the hulls being a fibrous component with little or no perceived nutritional value normally are separated and removed by physical means. The hypocotyls become separated following the splitting of the cotyledons, and while these currently generally are not deliberately isolated, they may be separated and isolated by passing the disturbed soyabeans over a sieve of sufficient pore size to selectively remove the small hypocotyl. The hypocotyl contains approx. 1.0-1.5 g% isoflavones (95% daidzein, 5% genistein). The raw hypocotyl and hull material can be ground or milled to produce, for example, a dry powder or flour which then could be either blended or used separately as a dietary supplement in a variety of

ways including, for example, as a powder, in a liquid form, in a granulated form, in a tablet or encapsulated form, or added to other prepared foodstuffs. Alternatively, it could be further processed to yield an enriched extract of phyto-oestrogens. Either or both of these materials also could be added to other leguminous material such as clover to provide the invention.

In plants, the oestrogenic isoflavones are restricted principally to the leaf, fruit and root; the stem and petiole contain very little. With soya and other common human legume foodstuff crops, the leaves are rarely regarded as foodstuff; indeed with these crops, the plants normally are allowed to die and dry out before the seed crop is harvested. Nevertheless, the fresh leaves of these crops could be regarded as a source of phyto-oestrogens for the invention although the much lower isoflavone content of the leaves of these crops compared to clovers, plus their generally slow growth compared to clovers, suggests that they would not be a preferred source of large-scale isoflavone enrichment.

To provide a similar amount of isoflavone to that contained in most traditional legume-rich diets (50-100 mg oestrogenic isoflavones/day) would require an average daily consumption of 3-6 g dry weight or 15-30 g wet weight of specially selected cultivars of clover with particularly high isoflavone levels. Clover grasses generally are not eaten by humans, except to a limited extent as sprouts of some of the pleasanter tasting varieties. Isoflavones are intensely astringent and are responsible in large part for the bitter taste of legumes. Thus the types of bean sprouts, clover sprouts and alfalfa sprouts generally available have been selected on the basis of cultivar and of age for pleasant taste, and in so doing inadvertently have been selected for low isoflavone content. Of the sprouts currently available in Western countries for human consumption, between approx. 100-250 g would need to be consumed daily to provide a dosage of 50-100 mg isoflavones. Certainly clovers and other legume sprouts are not generally eaten in such sufficient quantities by humans to obtain the advantages of the present invention.

The invention also concerns formulations containing the phyto-oestrogens discussed above together with a dietary suitable excipient, diluent, carrier, or with a food. Ideally the formulation is in the form of a pill, tablet, capsule, or similar dosage form.

The formulations may be a variety of kinds, such as nutritional supplements, pharmaceutical

preparations, vitamin supplements, food additives or foods supplemented with the specified active phyto-oestrogens of the invention, liquid or solid preparations, including drinks, sterile injectable solutions, tablets, coated tablets, capsules, powders, drops, suspensions, or syrups, ointments, lotions, creams, pastes, gels, or the like. The formulations may be in convenient dosage forms, and may also include other active ingredients, and/or may contain conventional excipients, carriers and diluents. The inclusion of the subject phyto-oestrogens in herbal remedies and treatments is also a preferred part of the invention.

The invention is also directed to the amelioration, prevention, or of various conditions responsive to treatment with the phyto-oestrogen substances of the invention. The preferred amounts to be administered to the human fall within 20 - 200 mg on a daily basis. More preferably the dosage is from 50 - 150 mg on a daily basis, and most preferably at a dosage of about 100 mg. If desired greater dosages can be administered for therapeutic reasons. In contrast to prior practices such high dosages were not possible. For example, dosages of up to or greater than 1000 mg may be suitable to treat some conditions. In order to obtain the benefits of the invention, the treatment with the isoflavones should continue for a considerable period, ideally for at least a month, and ideally continuously for the whole period for which the health improvement advantages should accrue.

The product according to the present invention yields a constant and accurately known amount of isoflavones. The product is also ideally a natural product, which has advantages for consumer acceptance, and in accordance with the supposed theory behind the invention may very possibly be one of the main causes for its beneficial effects. Whole legumes have a widely variable isoflavone content due to two main causes: the type of legume and the environmental effect. The type of legume typically has a wide range of isoflavone content. The miligram of isoflavone per hundred gram of whole foodstuff (dry weight) is given in the following table:

## Soya Products

T160X

- Whole Soya	150 - 300
- Soya Milk	25 - 40 (mg per 200 ml)
- Tofu	55 - 95
Lentils	80 - 120
Chickpeas	70 - 130
Broad beans	15 - 20
Garden peas	15 - 25

Thus common leguminous foodstuffs consumed in Western countries (broad beans, garden peas etc) have relatively low oestrogenic isoflavone content and exceptionally large amounts of these would need to be consumed daily to approximate those isoflavone levels consumed in traditional diets. Most Western cultures do not traditionally eat legumes with high isoflavone contents, and those soya products (milk, tofu etc) which are becoming increasingly popular in Western countries, also have relatively low isoflavone levels compared to whole soya, indicating that relatively large amounts of these would need to be consumed on a regular basis to deliver the required isoflavone levels.

The enviromental effect arises because the isoflavone levels in any species of plant depend greatly on the age of the plant, the climatic conditions where it is grown, the fertiliser and so forth. Therefore constant and consistant dosage is very difficult with ordinary whole foodstuffs. The accurately determined quality and quantity of the active isoflavones in the product, and its easy consumability when compared with the almost impossible task of eating huge amounts of often practically inedible foods, is therefore an import feature of the invention for preventing and helping in overcoming various health problems.

Among the various health problems, the treatment or prevention of high blood cholesterol levels, and the treatment of PMS and menopausal symptoms is especially important.

The product of the invention modulates the production and/or function of endogenous sex hormones in humans to modify or produce health improving effects, including the following:

- (i) lowered levels of various blood lipoproteins including, for instance, low-density and very-low-density cholesterol leading to reduced risk of development of atherosclerosis;
- (ii) reduced risk of development of cancer of the prostate;
- (iii) reduced risk of cancer of the

breast; (iv) reduced risk of development of cancer of the uterus; (v) reduced risk of development of cancer of the large bowel; (vi) reduced risk of development of the syndrome in women commonly referred to pre-menstrual syndrome (PMS), which includes pre-menstrual tension (PMT); (vii) reduced risk of development of many untoward symptoms (including dry vagina, peripheral flushing, depression etc) commonly associated in women with menopause; and for treating benign breast disease in women (benign or cystic breast disease associated with non-malignant swelling and tenderness of breast tissue). The invention therefore is directed to a method for the prophylaxis or treatment of a human, to combat conditions associated with phyto-oestrogen deficiency, which comprises administering to the human an effective amount of phyto-oestrogen principally isoflavone but which might also include relatively smaller amounts of lignans and coumestans, ideally in a concentrated form, wherein the isoflavones include genistein, and/or biochanin A, and/or daidzein, and/or formononetin.

Cancer of the breast generally is considered to be associated with oestrogenic dysfunction. Breast cancer cells may display more oestrogen receptors than normal breast cells and stimulation of these receptors by endogenous oestrogens is thought to be a prime source of stimulation of their malignant growth. Currently synthetic anti-oestrogens are being used to prevent or treat the growth of malignant breast cells. Isoflavones are potent anti-oestrogens that could be expected to help prevent or to successfully treat breast cancer. It has been reported that the risk of breast cancer in western societies is indirectly proportional to the level of phyto-oestrogens in the diet and to the amounts of phyto-oestrogen metabolites excreted in the urine.

Cancer of the prostate generally is considered to be associated with sex hormone dysfunction and the growth of prostatic cancer cells is influenced by oestrogens and androgens. The incidence of prostatic cancer is low in communities with high legume intake and, conversely, is high in Western societies. Phyto-oestrogens are thought to protect from development of prostatic cancer. One mechanism may be the effect of phyto-oestrogens on lowering the proportion of unbound reproductive hormones in the blood. However, there is other evidence to suggest that phyto-oestrogens, particularly isoflavones, can have a direct influence on certain cellular enzymes within prostatic cells.

Pre-menstrual syndrome has uncertain aetiology and pathogenesis, although most certainly is

associated with reproductive hormone dysfunction. It also is a syndrome which has reportedly lower incidence in communities maintaining traditional high-legume diets. It is proposed that phyto-oestrogens will alleviate this condition by restoring balance to oestrogen metabolism.

Menopausal syndrome is associated with changes in the oestrogen profile in the body with advancing age. Adverse clinical symptoms may be treated with oestrogen replacement therapy. There is evidence that foodstuffs high in phyto-oestrogens are a suitable alternative to synthetic hormones in this respect, producing alleviation of adverse clinical symptoms. Again, it is proposed that phyto-oestrogens will function by restoring balance to oestrogen metabolism.

Benign (or cystic) breast disease has unknown aetiology. However, its association in women with certain stages of the menstrual cycle is strongly suggestive of oestrogen dysfunction. There currently is no successful treatment of this condition. Phyto-oestrogens are proposed to successfully treat this condition by restoring balance to oestrogen metabolism.

Atherosclerosis is associated with cholesterol metabolism which in turn is associated closely with oestrogen metabolism. The generally higher incidence of atherosclerosis in young men versus young women, the rising incidence in women following menopause, and the lower incidence in post-menopausal women receiving oestrogen replacement therapy, all point to the moderating influence of oestrogens on cholesterol metabolism. A prime effect of oestrogens on cholesterol metabolism is stimulation of the liver to process cholesterol, particularly the highly atherogenic low-density lipoproteins and very low-density lipoproteins, into bile salts. It is proposed that phyto-oestrogens have an important hypocholesterolaemic effect in humans. There may be a variety of mechanisms involved, but one may be the stimulation by phyto-oestrogens of cholesterol catabolism by the liver.

#### MODES FOR CARRYING OUT THE INVENTION

The invention is now described with reference to various examples.

##### EXAMPLE 1 - Preparation of Red Clover Product

Tablets were prepared using red clover in accordance with the following procedure. The raw plant material is harvested and dried; such drying being either sun-drying or from applied heat.

18

The dried product is then preferably chaffed, before the following extraction step, although this can be omitted if desired.

The dried material is extracted in an aqueous: organic solvent mix. The aqueous phase is required to extract the water-soluble glycoside form of isoflavones, while the organic solvent is required to solubilise the water-insoluble aglycone form. The organic solvent can be either alcohol, chloroform, acetone or ethyl acetate. The ratio of solvent in the water can be between 0.1% and 99.9%. The preferred method is to use 60% alcohol in water.

The isoflavones are extracted by exposing the plant material to the water:solvent mix. The exposure time in general terms is indirectly proportional to the temperature of the mixture. The temperature of the mix can range between ambient temperature and boiling temperature. The exposure time can be between 1 hour and 4 weeks or even longer. It has been determined that the adequate times for maximal recovery of isoflavones are 2 weeks at 50°C and 24 hours at 90°C. The supernatant is separated from the undissolved plant material and the organic solvent removed by distillation. The aqueous supernatant then is concentrated, typically by distillation.

Additional processing steps can be used, if desired, to convert the extracted natural product to capsule, tablet, or other convenient form for ingestion, using normal techniques for doing this. Otherwise the product can be packaged as a convenient food additive.

#### EXAMPLE 2 - preparation of soya hypocotyl product.

Soyabeans were heated in dry air so that the hull became brittle. The beans then were processed through a tumble mill which removed the hull and split the bean the two cotyledons and the small-sized hypocotyl which separated from each other. The light-weight hulls then were removed by an air stream. The small-sized hypocotyls were separated from the larger cotyledons by sieving through a steel wire mesh with apertures of 1 mm x 1 mm. This yielded approximately 87% purity of hypocotyls with 13% contamination by small cotyledon chips.

Normal soybean processing steps isolate the hulls and then these are discarded or processed

19

separately for use in human and animal foodstuffs. The hypocotyls normally are not separated and are processed along with the cotyledons. However, a small number of soybean processors are separating hypocotyls by the above methods in order to reduce the astringent taste of soyflour for human consumption, and currently these hypocotyls are either discarded or processed to flour for use in animal feed.

#### EXAMPLE 3 - effect of administering red clover extract to humans

Seven normal individuals were studied for the comparative effects of red clover extract and whole legumes on blood cholesterol levels. All the individuals were consuming a standard Western diet with minimal levels of legumes.

Three men consumed between 100-150 g haricot or navy beans daily for 3 weeks as a supplement to their normal diet. This yielded an approximate daily isoflavone dosage or between 60-100 mg.

Four other individuals (3 men, 1 woman) consumed 5 g of red clover extract containing 100 mg isoflavones daily for 3 weeks.

Total serum cholesterol levels were determined immediately before and immediately following the challenge.

	<u>Pre-treatment</u>	<u>Post-treatment</u>	<u>% change</u>
<u>Beans only</u>			
T200X Patient 1	5.77	5.46	- 5.4
Patient 2	6.24	6.12	- 1.9
Patient 3	7.45	8.51	+14.3
<u>Red clover extract</u>			
Patient 5	6.53	5.90	- 9.6
Patient 6	7.43	6.63	-10.8
Patient 7	6.33	5.50	-13.1
Patient 8	6.98	7.28	+ 4.3

The red clover extract had a significantly ( $P < 0.05$ ) greater hypocholesterolaemic effect than



did the whole beans.

Neither of the treatments produced any untoward side effects, although the whole bean eaters reported greater difficulty with compliance of treatment than did those taking the red clover extract.

#### EXAMPLE 4 - effect of administering soy hypocotyls to humans

Fifteen volunteers (8 women, 7 men) were given 5 g of soy hypocotyl containing (45 mg daidzein and 5 mg genistein) daily for 2 months. The hypocotyl was consumed as a powder added to the diet.

The effects on cholesterol levels are shown in the following table. The individuals are grouped according to their pre-treatment cholesterol levels (high, medium, low).

T219x

		Range (mean) unmol/L	
	n	Pre-treatment	Post-treatment
Group 1	6	6.3 - 8.4 (7.1)	5.4 - 6.5 (6.1)
Group 2	6	5.0 - 6.2 (5.5)	4.7 - 5.9 (5.1)
Group 3	3	3.3 - 4.7 (4.2)	3.4 - 4.6 (4.1)

The results show a significant fall in total cholesterol levels in those individuals with cholesterol levels considered to be at the upper end of the normal range.

In addition, 1 woman reported substantial amelioration of her benign breast disease problem associated with mid-cycle swelling and tenderness, and another woman reported regularisation of her menstrual cycle and reduced menstrual bleeding. Both of these effects were regarded as beneficial.

No other side-effects were reported as a result of the treatment.

THE CLAIMS:

1. A health supplement comprising a health supplementary amount of a phyto-oestrogen selected from genistein, daidzein, biochanin A, and/or formononetin.
2. The supplement according to claim 1 which also comprises at least one dietary suitable excipient, diluent, carrier or food.
3. The supplement according to claim 1 wherein said phyto-oestrogen is extracted from red clover.
4. The supplement according to claim 1 wherein said phyto-oestrogen is extracted from soya.
5. The supplement according to claim 4 wherein said phyto-oestrogen is extracted from soya hypocotyls.
6. The supplement according to claim 1 wherein said phyto-oestrogen comprises genistein, and/or biochanin A : daidzein and/or formononetin, present in a ratio of from about 1:2 to 2:1.
7. The supplement according to claim 1 in unit dosage form, wherein said phyto-oestrogen is present in an amount of from about 20 mg to 200 mg per dosage unit.
8. The supplement according to claim 7, where the amount is 50 to 150 mg.
9. The supplement according to any one of claims 1 to 8, which is in the form of a tablet or capsule.
10. A method of improving the health of a human which comprises administering to said human a health supplementing amount of a phyto-oestrogen selected from genistein, daidzein, biochanin A, and/or formononetin.

22

11. The method according to claim 10, wherein said phyto-oestrogen is extracted from red clover.

12. The method according to claim 10, wherein said phyto-oestrogen is extracted from soy.

13. The method according to claim 12, wherein said phyto-oestrogen is extracted from soy hypocotyls.

14. The method according to claim 10, whereby said phyto-oestrogen comprises genistein and/or biochanin A : daidzein and/or formononetin, present in ratio of from about 1:2 to 2:1.

15. The method according to claim 10, wherein the phyto-oestrogen is administered in an amount of from about 20 mg to 200 mg per day.

16. The method according to claim 10, wherein the phyto-oestrogen is administered in an amount of from about 50 mg to 150 mg per day.

17. The method according to claim 10 whereby the phyto-oestrogen is administered at least daily, over a period of at least a month.

18. The method according to any one of claims 10 to 17 for improving the health of a human female who has, or may develop, a condition selected from any one or more of: breast cancer, benign breast disease, pre-menstrual syndrome, or symptoms associated with menopause.

19. The method according to ~~any one of claims 10 to 17~~ <sup>claim 10</sup> for improving the health of a human who has, or may develop, elevated levels of cholesterol in the blood stream.

20. The method according to any one of claims 10 to 17 for improving the health of a human who has or may develop cancer.

23

add  
C77

add  
C17

## AMENDED CLAIMS

[received by the International Bureau on 8 October 1993 (08.10.93);  
original claims 1-10 amended; other claims unchanged (1 page)]

1. A health supplement comprising a health supplementary amount of a naturally occurring phyto-oestrogen selected from any two or more of genistein, daidzein, biochanin A, formononetin, and/or their glycosides.
2. The supplement according to claim 1 which also comprises at least one dietary suitable excipient, diluent, carrier or food.
3. The supplement according to claim 1 wherein said phyto-oestrogen is extracted from red clover.
4. The supplement according to claim 1 wherein said phyto-oestrogen is extracted from soya.
5. *composition* The ~~supplement~~ according to claim 4 wherein said phyto-oestrogen is extracted from soya hypocotyls.
6. The supplement according to claim 1 wherein said ~~phyto-oestrogen~~ comprises genistein, and/or biochanin A : ~~daidzein and/or formononetin~~, present in a ratio of from about 1:2 to 2:1.
7. The supplement according to claim 1 in unit dosage form, wherein said phyto-oestrogen is present in an amount of from about 20 mg to 200 mg per dosage unit.
8. *composition* The ~~supplement~~ according to claim 7, where the amount is 50 to 150 mg.
9. The supplement according to ~~any one of claims 1 to 8~~, which is in the form of a tablet or capsule.
10. A method of improving the health of a human which comprises administering to said human a health supplementing amount of a naturally occurring phyto-oestrogen selected from any two or more of genistein, daidzein, biochanin A, formononetin, and/or their glycosides.

STATEMENT UNDER ARTICLE 19

The independent claims have been restricted to define a supplement containing at least two of the naturally occurring phyto-oestrogens, including their glycosidic forms.